

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|--|
| (51) International Patent Classification ⁵ : A01N 43/16, 25/26, 25/24 A01C 1/06 | A1 | (11) International Publication Number: WO 93/25078 (43) International Publication Date: 23 December 1993 (23.12.93) |
| (21) International Application Number: PCT/FI93/00262 (22) International Filing Date: 16 June 1993 (16.06.93) (30) Priority data: 922780 16 June 1992 (16.06.92) FI (71) Applicant (for all designated States except US): OY ALKO AB [FI/FI]; Salmisaarenranta 7, FIN-00180 Helsinki (FI). (72) Inventor; and (75) Inventor/Applicant (for US only) : KOISTINEN, Tarja [FI/FI]; Toivojentie 6, FIN-05200 Rajamäki (FI). (74) Agent: LÖNNQVIST, Gunnel; Oy Alko Ab, Law Department/Patents, P.O. Box 350, FIN-00101 Helsinki (FI). | | (81) Designated States: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> |
| (54) Title: COMPOSITIONS AND METHODS FOR COATING SEED (57) Abstract <p>The present invention is related to a seed coating which is beneficial for the germination of seeds. A polysaccharide derivative hydrolysate, especially a hydrolysate of carboxymethyl cellulose (CMC) or carboxymethyl starch (CMS), is used as a binder, which is capable of binding pesticides and herbicides and talcum as a thin, even and unspotty film. In the method according to the invention the binder can be sprayed on the seed as a concentrated water solution, preferably a 30 % (w/v) solution.</p> | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|--|----|--------------------------|
| AT | Austria | FR | France | MR | Mauritania |
| AU | Australia | GA | Gabon | MW | Malawi |
| BB | Barbados | GB | United Kingdom | NL | Netherlands |
| BE | Belgium | GN | Guinea | NO | Norway |
| BF | Burkina Faso | GR | Greece | NZ | New Zealand |
| BG | Bulgaria | HU | Hungary | PL | Poland |
| BJ | Benin | IE | Ireland | PT | Portugal |
| BR | Brazil | IT | Italy | RO | Romania |
| CA | Canada | JP | Japan | RU | Russian Federation |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan |
| CG | Congo | KR | Republic of Korea | SE | Sweden |
| CH | Switzerland | KZ | Kazakhstan | SK | Slovak Republic |
| CI | Côte d'Ivoire | LI | Liechtenstein | SN | Senegal |
| CM | Cameroon | LK | Sri Lanka | SU | Soviet Union |
| CS | Czechoslovakia | LU | Luxembourg | TD | Chad |
| CZ | Czech Republic | MC | Monaco | TG | Togo |
| DE | Germany | MG | Madagascar | UA | Ukraine |
| DK | Denmark | ML | Mali | US | United States of America |
| ES | Spain | MN | Mongolia | VN | Viet Nam |
| FI | Finland | | | | |

COMPOSITIONS AND METHODS FOR COATING SEED

5

THE BACKGROUND OF THE INVENTION

1. The Field of the Invention

10

The present invention is related to a seed coating composition and a method for coating seeds with polysaccharide derivative hydrolysates.

2. The State of the Art

15

Seeds are coated to improve germination, to retain humidity and bind pesticides, herbicides and fertilizers.

20

Commonly used coatings are polyacrylates (W085/01736) and different starch compositions (US 4,250,660). In the patent US 4,250,660 a 3-5 % binder solution of unhydrolysed carboxymethyl cellulose (CMC) is described.

25

Unhydrolysed polysaccharide derivatives have a relatively low solubility in water. CMC is the most water soluble of the unhydrolysed polysaccharide derivatives. In spite of that, it is an acknowledged fact that, when the amount of unhydrolysed polysaccharide derivatives, especially CMC, is raised, the viscosity of the solution increases rapidly. The increase of viscosity makes the handling of the solution difficult, e.g. pumping or spraying of the solution is impossible. A dilute solution of unhydrolysed CMC on the other hand is not capable of binding the coating strongly enough.

30

35

To improve the feasibility of unhydrolysed CMC in seed coating, CMC has been combined with other substances. Unhydrolysed CMC has been used as very dilute solutions (0,5 - 1 %) for coating cotton seeds in combination with lignin (SU 1,015,837).

When seeds are coated with unhydrolysed polysaccharide derivatives, cleaning of the equipment is very tedious. Toxic wastes containing pesticides and herbicides are produced. The destruction of such hazardous wastes always causes problems.

The present invention is based on experiments performed with polysaccharide derivative hydrolysates, especially CMC and CMS-hydrolysates. The results indicate that the solubility and viscosity of polysaccharide derivatives differ greatly. We have observed that a hydrolysed CMC can be pumped as a 40 % (w/v) solution, whereas the most water-soluble of the polysaccharides, CMC, and more specifically a brand of CMC, which has an especially low viscosity, can be pumped only when the concentration of the solution is less than 10 % (w/v).

Said observation indicates that polysaccharide derivative hydrolysates, preferably CMC- and CMS-hydrolysates, are extremely suitable for coating seeds, especially seeds from oleiferous plants.

THE SUMMARY OF THE INVENTION

The present invention provides a seed coating composition, which as a binder uses polysaccharide derivate hydrolysates, preferably CMC- or CMS-hydrolysates, which can be used as sufficiently concentrated solutions. The viscosity of the solutions is still so low that solutions, which can be pumped or sprayed, are easy to prepare and handle.

Particularly, the present invention provides a new method, in which the coating does not adhere to the walls of the coating equipment and the solution, which has been sprayed on the walls, is easy to rinse away with water alone.

The method is especially suitable for coating seeds from oleiferous plants. Seeds from such plants are known to be difficult to coat, because of their extremely oily and slippery surface.

5

The present invention provides a coating, which comprises a tight, non-spotty, durable film, which decomposes easily in the soil.

10

The present invention also provides a coating, which neither adheres to the equipment nor produces seed clusters. The coated seeds should preferably be separate. It is also important that the germination is not disturbed by the coating or binder.

15

Furthermore, the present invention provides a binder, which is capable of binding pesticides, herbicides, nutrients, fertilizers or microbes as well as talcum to the surface of the seed as a thin, durable, unspotty film.

20

Polysaccharide derivative hydrolysates, especially CMC-hydrolysates but also CMS-hydrolysates, which can be used as sufficiently concentrated solutions, e.g 5 - 50, preferably 10 - 40, most preferably 30 % (w/v) water solutions are suitable binders for seed coatings according to the present invention.

25

30

The cellulose or starch derivative hydrolysates according to the present invention should have an average degree of polymerisation in the range of 3 - 500, preferably 5 - 300, most preferably 5 - 100 and they are optionally substituted by carboxymethyl, methyl, methylethyl, hydroxypropyl, hydroxypropylmethyl or hydroxyethyl groups.

35

It is important that the polysaccharide derivative hydrolysate used in the method of the present invention can be pumped or sprayed on the seeds.

THE DETAILED DESCRIPTION OF THE INVENTION

5 The polysaccharide derivative hydrolysates used as coatings
for seeds are preferably hydrolysates of cellulose or
starch derivatives. The average degree of polymerisation
of the hydrolysates is in the range of 3 - 500, preferably
5 - 300, most preferably 5 - 100. The cellulose or starch
10 is preferably substituted by carboxymethyl, methyl,
hydroxypropyl, hydroxypropylmethyl or hydroxyethyl groups.
The properties and preparation of said hydrolysates of
polysaccharide derivatives used as seed coatings are
described in more detail in the following patent
applications EP 382 577, EP 382 576, EP 382 578,
15 EP 470 855, EP 470 870, EP 470 872, EP 470 871 and
US 07/743,152.

Such preparations are produced preferably by enzymatic
degradation from substantially purified polysaccharide
20 derivative solutions. These solutions are relatively di-
lute. Because the product is dried after the degradation,
very concentrated water solutions, containing up to 40 %
(w/v) of polysaccharide derivative hydrolysate in water can
be produced. These solutions are easy to handle because
25 they have a relatively low viscosity in spite of the
concentration.

The polymerisation degree or molecular weight as well as
concentrations and viscosities for some representative
30 starting materials for the polysaccharide derivative hydro-
lysates as well as the polysaccharide derivative hydro-
lysates are shown in the Table 1.

The viscosities were determined with a Haake viscosimeter.
35 The viscosity of a 30 % water solution of unhydrolysed CMC
and CMS could not be measured. The viscosity of a 2 %
water solution of hydrolysed CMC or CMS was negligible.

Table 1

| POLYSACCHARIDE DERIVATIVE | VISCOSITY/CONCENTRATION | |
|---------------------------|-------------------------|---------------|
| | UNHYDROLYSED | HYDROLYSED |
| CMC | 12 mPas/ 2 % | 34 mPas/ 30 % |
| CMS | 5 mPas/ 2 % | 20 mPas/ 30 % |

Seeds from e.g tomato, wheat, oats, oleiferous plants, soft wood (conifers), beets, maize, cotton and vegetables are coated with a thin film of a composition containing binders, active components and fillers. The coating is used to improve germination by increasing the retention of moist. A good capability of binding pesticides, herbicides, fertilizers, nutritions and advantageous microbes also improves the quality of the seeds. The method of the invention is especially suitable for coating oleiferous plants.

In coating experiments it was observed that polysaccharide derivative hydrolysates, especially CMC-hydrolysate and CMS-hydrolysates did not adhere to the surface of the the equipment. In addition, the solution sprayed on the walls could easily be rinsed away with water.

In the method according to the invention hydrolysates of the polysaccharide derivatives can be used as solutions of 5 - 50, preferably 10 - 40, most preferably 30 % (w/v) concentration.

The seed coating can be performed in mixing drums. The seeds are coated with hydrolysates of polysaccharide derivatives, preferably hydrolysates of CMC or CMS. The hydrolysates of the polysaccharide derivative form on the sur-

face of the the seed a thin film. The pesticides, herbicides, fertilizers and/or talcum adheres evenly to the thin film of the binder. It is essential that the binder does not cause the seed to form clusters. The seeds should roll
5 freely in the drum without adhering to each others or the walls of the mixer.

The coating can also be performed in a coating equipment, which preferably has a screw driver (conveyor), on which
10 the seeds to be treated are portioned out. On the seeds different binders and pesticides, herbicides, nutritions, fertilizers and talcum are sprayed. In seed coating, it is important that the binder remains moist until the talcum and the active ingredients have been portioned out on the
15 surface of the seed. In seed coating the film should be even, tight, non-spotty and durable. At the same time it should be easily dissolvable in soil.

The invention is described in more detail in the following
20 examples, in which some characteristic modes of seed coating with some conventional binders suitable for coating seeds are compared with the binder of the present invention. The examples are meant to illustrate the invention without restricting the scope of the the invention
25 in any way.

Example 1

CMC-hydrolysates (CMC-H)

a) Tests using CMC-H as a coating

30 In a mixing drum seed of oleiferous plants were coated in 1,5 kg portions with CMC-H solutions of different concentrations in order to find the best CMC-H concentration for coating. CMC-H forms a uniform, thin film on the surface of the seed. The seeds do not adhere to each others
35 or the equipment. The seeds roll in the drum without adhering to each others or the walls of the mixer. The 30 % (w/v) solution proved to be the best.

b) Tests using CMC-H as a binder

5 The coating equipment consisted of a 7 meter screw driver,
whereon seeds were portioned out. Different binders such
as CMC-H solutions and polyacrylates were compared and
sprayed on the seeds. The best concentration of CMC-H was
30 % (w/v). The final percentage of the dry binder per
kilogram of seed was 4 % (w/w) CMC-H. Depending upon the
10 binder 2 % pesticide and 2 - 3 % talcum was added per kilo-
gram seed.

15 The seeds coated with CMC-H remained moist until the talcum
was portioned out. The talcum formed a thinner and more
even film when CMC-H was used. When polyacrylate was used,
the film was much more spotty. In other words the film was
not even and it had holes in the film layer.

20 CMC-H did not form seed clusters and did not adhere to the
equipment. In the method no secondary products, such as
clusters or debris, containing pesticides or herbicides
were produced. Thus, no hazardous wastes had to be disposed
of....

Example 2

25 Test using polyacrylate (Uusi Kiilto) as a binder.
The test was performed as described in Example 1b using a
27 % (w/v) acrylate solution (Uusi Kiilto). The amount of
dry binder per kilogram of seed was 4 - 6 % (w/w).

30 The binder adhered to the equipment and formed clusters of
seed. The equipment had to be cleaned using strong mecha-
nical force. The film of the coating was spotty.

35 The clusters, which could not be broken up, caused losses
in many ways. The losses were caused partly, by the ad-
herance to the equipment and partly, because the clustered
seeds were not first class products, which could be used

and sold. Furthermore, the wastes caused problems, because they had to be treated as hazardous wastes. In addition, dust classified as hazardous was released when the seeds were screened to separate the free seeds from the clusters.

5

Example 3

Test using polyacrylate (Hilleshög) as a binder

The test was performed as described in Example 1b using a polyacrylate (Hilleshög) solution having a 30 % (w/v) concentration. The content of dry binder per kilogram of seed was 4 - 6 % (w/w). The results corresponds to those described in Example 2.

10

Example 4

15 CM-Starch hydrolysate (CMS-H)

a) Test using CMS-H as a coating

In a mixing drum seeds of oleiferous plants (Kesko 0303 - hard) were coated with solutions of CMS-hydrolysate as described in Example 1. The purpose of the test was to find the best CMS-H concentration for seed coating. In the test three somewhat different types of solutions of CMS-H (PRIMOJEL-K, CMC-H 24.28 and CMS 30.34) were used. With all these preparations similar results were achieved. In all tests the 30 % (w/v) CMS-H solution proved to be the best.

20

25

b) Test using CMS-H as a binder

The coating device was a 7 meter long screw driver, on which the seeds were portioned out. A water solution containing 30 % (w/v) CMS-H described in Example 4a was sprayed on the seeds, as well as 2 % pesticide and 3 % talcum.

30

35

The structure of the coating was good and even. The pesticide and talcum added was bound excellently by the binder. CMS-H did not form clusters and it did not adhere to the surface of the equipment.

Example 5

Test with unhydrolysed CMC as a coating.

5 In a mixing drum 0,5 kg of seeds from oleifereous plants (Kesko 0303 - hard) was coated with with a CMC solution as described in Example 1 in order to find the best CMC concentration for the coating. In the test the CMC quality with the lowest possible viscosity (Finnfix 2) was used.

10 Using a 10 % (w/v) CMC-solution the coating capacity of the binder disappeared in about 3 minutes from the spraying. The reason was that the drying had started. CMC could bind neither the pesticide nor the talcum.

15 A 15 % (w/v) CMC-solution was so thick that it was impossible to spray it as a mist on the surface of the seeds. The CMC-solution formed droplets on the surface and clusters were formed when the talcum was added.

20 A suitable CMC concentration was not found, because the 10 % (w/v) solution was too dilute for binding and the 15 % (w/v) solution was too viscose for spraying.

Example 6

25 Germination test

The germination test was done according to standardised prescriptions (directives) issued by the Finnish State Institute for Seed Control. The seeds were germinated on blotting paper for 3 weeks in 10 °C and whereafter for 7
30 days in 20 °C.

The seeds were a hard brand of rape (Kesko 0303). The results are shown in Table 2.

35

Table 2

| | Coating | Germinability |
|----|--|---------------|
| 5 | Uncoated | 94 |
| | Binder (27 % (w/v) solution), Uusi Kiilto, 4-6 w/v % binder/kg seed | 96 |
| 10 | Binder (27 % (w/v) solution), Uusi Kiilto 6-8 w/v % binder/kg seed | 93 |
| | Binder (30 % (w/v) solution), Hilleshög 4-6 % (w/v) binder/kg seed | 97 |
| 15 | Binder (30 % (w/v) solution), CMC-H 4 % (w/v) binder/kg seed | 95 |

The differences in germinability were not significant and
the binders did not disturb germinability.

25

30

35

WHAT IS CLAIMED IS:

1. A seed coating composition comprising as a binder a
5 5 - 50, preferably 10 - 40, most preferably 30 % (w/v)
water solution of a polysaccharide derivative hydrolysate,
preferably a hydrolysate of a cellulose or starch derivati-
ve having an average degree of polymerisation in the range
of 3 - 500, preferably 5 - 300, most preferably 5 - 100.
- 10 2. The seed coating composition of claim 1, wherein the
cellulose or starch derivative hydrolysate is optionally
substituted by carboxymethyl, methyl, methylethyl, hydroxy-
propyl, hydroxypropylmethyl or hydroxyethyl groups.
- 15 3. The seed coating composition of claim 3, wherein the
cellulose or starch derivative hydrolysate is CMC or CMS
hydrolysate.
- 20 4. A method for coating seeds, wherein the seeds are trea-
ted with a binder comprising a 5 - 50, preferably 10 - 40,
most preferably 30 % (w/v) water solution of a poly-
saccharide derivative hydrolysate, preferably a hydro-
lysate of a cellulose or starch derivative having an ave-
25 rage degree of polymerisation in the range of 3 - 500,
preferably 5 - 300, most preferably 5 - 100.

5. The method of claim 4, wherein the cellulose or starch derivative hydrolysate is a CMC or CMS hydrolysate.
6. The method of claim 4, wherein the seeds are treated by spraying.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI '93/00262

A. CLASSIFICATION OF SUBJECT MATTER

IPC5: A01N 43/16, A01N 25/26, A01N 25/24, A01C 1/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC5: A01N, A01C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| X | DE, A1, 2920568 (CHINOIN GYOGYSZER ES VEGYESZETI TERMEKEK GYARA RT), 29 November 1979 (29.11.79), page 3, line 1 - page 6, line 31, claims 1-18 --- | 1-6 |
| X | Plant Disease, Volume 69, No 12, December 1985, A. B. Filonow et al., "Evaluation of Several Actinomycetes and the Fungus Hyphochytrium catenoides as Biocontrol Agents for Phytophthora Root Rot of Soybean" page 1033 - page 1036 --- | 1-6 |
| X | WO, A1, 9000351 (AQUASPERSIONS LIMITED ET AL.), 25 January 1990 (25.01.90), claims 1-7, page 2, line 14-line 31; page 3, line 20-line 25; page 5, line 1-line 4; page 9, line 14-line 16, line 24-line 32 --- | 1-6 |

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 Sept 1993

Date of mailing of the international search report

14 -09- 1993

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Gerd Wranne
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 93/00262

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| A | EP, A1, 0382576 (ALKO LTD), 16 August 1990 (16.08.90), page 5, line 36 - line 40 -- | 1-6 |
| A | EP, A1, 0470872 (ALKO LTD), 12 February 1992 (12.02.92) -- | 1-6 |
| A | DE, C, 921291 (AGRICULTURAL RESEARCH CORPORATION), 4 November 1954 (04.11.54), page 3, line 41 - line 51, claims 1,2,6,7 -- | 1-6 |
| A | DE, C, 964736 ("BIO" NÄHRMITTELFABRIK DR. SCHWEIGMANN & CO.), 29 May 1957 (29.05.57) -- ----- | 1-6 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

30/07/93

International application No.

PCT/FI 93/00262

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|--|--|
| DE-A1- 2920568 | 29/11/79 | AT-B- 366555 BE-A- 876494 CA-A- 1118221 FR-A,B- 2426412 GB-A,B- 2022568 JP-A- 55017370 JP-B- 61057801 NL-A- 7904016 SU-A- 1246876 US-A- 4272276 | 26/04/82 17/09/79 16/02/82 21/12/79 19/12/79 06/02/80 09/12/86 27/11/79 23/07/86 09/06/81 |
| WO-A1- 9000351 | 25/01/90 | AU-B- 633504 AU-A- 3983589 EP-A- 0352010 EP-A- 0424454 | 04/02/93 05/02/90 24/01/90 02/05/91 |
| EP-A1- 0382576 | 16/08/90 | AU-B- 620207 AU-B- 631842 AU-A- 4933390 AU-A- 4933490 AU-A- 4933590 CA-A- 2009675 CA-A- 2009676 CA-A- 2009677 EP-A- 0382577 EP-A- 0382578 | 13/02/92 10/12/92 16/08/90 16/08/90 16/08/90 10/08/90 10/08/90 10/08/90 16/08/90 16/08/90 |
| EP-A1- 0470872 | 12/02/92 | AU-A- 8174491 CA-A- 2042529 JP-A- 4281836 | 13/02/92 11/02/92 07/10/92 |
| DE-C- 921291 | 04/11/54 | NONE | |
| DE-C- 964736 | 29/05/57 | NONE | |